BOOK REVIEW

By A. W. McCallum,

Forest Pathologist

Bagchee, K. Investigations on the infestation of *Peridermium himalayense*, Bagchee, on *Pinus longifolia*, Part II., *Cronartium himalayense*, n. sp., on *Swertia* spp. Distribution, Morphology of the Parasite, Pathological study of the Infection, Biological Relationship with the Pine Rust, and Control. Indian Forest Records 18 (11): 1-66. 17 pl. 1933.

A blister rust which attacks *Pinus longifolia* or chir—one of the pitch pines—has long been known to occur in north-western India. In recent years it has become epidemic, causing heavy losses in the young pine stands in the Kumaon and Garhwal Divisions. Trees up to 20 years old are most susceptible, while those older than 30 years seem to be immune. It is interesting to note that two American pitch pines—*Pinus arizonica* and *P. ponderosa*—growing in association with infected chir remained free from rust. Chir is one of the most important forest trees of the Himalayas.

In 1927 the alternate stage of this rust was discovered by the author, on *Swertia*—a small annual plant. Three species—*S. angustifolia*, *S. purpurescens*, and *S. alata*—were found in the field to be susceptible, and two species—*S. cordata* and *S. paniculata*—were found to be immune to rust. Artificial inoculations confirmed this observation. *Swertia* spp. are widely distributed on the slopes of the Himalayas from 3,000 to 10,000 feet in all forests where chir occurs. A complete technical description of the fungus is given and extensive details regarding the various spore forms. The method of infection of *Swertia*—by way of the stomata—and the development of the fungus in the alternate host are described in detail.

In regard to control, the author, after considering various possible methods, concludes that the eradication of the susceptible species of *Swertia* would be the only effective method. The inherent difficulties of this method are recognized but the belief is expressed that it would be worth while.

—A. W. M.