NEWS AND NOTES

FIELD NOTES

By I. C. M. Place

Prescribed Burning

Deliberate use of fire in forestry is not new on the continent. For years **Professor H. H. Chapman** (one of America's grand old men of forestry) preached its use to control hardwoods and prepare seedbeds for regeneration of loblolly and longleaf pine in the southern United States, much, it might be added, to the consternation of the U.S. Forest Service. His views are now regarded as orthodox. **Little** and **Moore** have developed prescribed burning to a fine art in New Jersey pine barrens. It has been used in jack pine in the Lake States and for years has been a standard method of reducing fire hazard and providing suitable seedbeds in logged off land in the West Coast. Although they were not interested in forestry, the Iroquois in northern New York state and the Hurons in southern Ontario used fire extensively to clear land for growing corn and the influence of these fires upon the forest has been considerable. Indeed, **Professor Armson** of the University of Toronto, has brought it to my attention that the great **Samuel de Champlain** made the humiliating confession in his journals that he lost his followers one day in an Indian corn field in southern Ontario.

In most of Canada, however, interest in prescribed burning has been largely academic and attempts to use fire have been sporadic and unsystematic. Perhaps one of the main reasons for this is that few people have had the courage to light the match. There are, however, signs of increasing interest. Several months ago, **Finn Frost**, Chief Forester for the Newfoundland Department of Mines and Resources, sent me a publication of the Newfoundland Research Committee devoted to a symposium on prescribed burning. The views are interesting and I would like to remark upon several of the papers.

Balsam fir is one of the most valuable pulpwood species in Newfoundland, and the balsam woolly aphid is currently threatening to lower drastically the productivity of Newfoundland's forests. Frost proposes that infested stands of fir be logged and burned both for sanitation and to prepare the land for either natural or artificial regeneration of black spruce which is not attacked by the aphid. His case is impressive.

D. E. Nickerson points out that the thriftiest pure stands of black spruce are invariably of fire origin and fire does indeed favor black spruce regeneration by reducing the dry surface layer of the seed bed. Seed, of course, is essential but studies by the Forestry Branch show the the serotinous cones at the top of ten to fifteen well distributed seed bearing trees will usually yield enough seed to reforest an acre.

H. W. Chancey, however cautions that constant severe burning over highly acid infertile soils in Newfoundland's humid climate is likely to intensify podsolization and iron pan formation. This could conceivably lead to a disastrous

drop in the permeability of the soil and extensive water logging and development of heathy barrens. The long-term effect of the black spruce itself on the soil would worry me. I have encountered apparently incipient hard pan formation in sandy soils under individual clumps of black spruce where it was lacking under adjacent trees of other species. Several other foresters have told me of similar experiences.

The whole question of prescribed burning is complex. Much depends on the individual circumstances, the species involved, the local climate and terrain and the objectives, whether these be seedbed preparation, control of brush or reduction of fire hazard. Obviously fire cannot be used on a hit or miss basis. Even where its utility is clearly apparent in certain forest types, one hesitates to recommend fire for indiscriminate use just as one would hesitate to recommend do-it-yourself X-ray treatment for cancer.

If prescribed burning is to be practicable, it must be applicable on moderately large areas at low cost. At present, information about the interactions of fire behavior, weather, and fuel moisture content in different cover types and the influence of these on control of burning is inadequate to say whether this is possible. Moreover, there are no data on the long term effects of fire on the soil for conditions prevailing in most of Canada. Considerable nitrogen is undoubtedly lost when the humus and litter on the forest floor are burned. Detailed studies in the southern Atlantic Coastal Plain have shown that this loss is quickly recouped there owing to nitrogen fixation by the leguminous plants that spring up after fire. Unfortunately, legumes are not abundant in our forests. Fire has been used most successfully in stands of intolerant trees, such as the pines, which grow on coarse soils. It is precisely these soils that can least stand depletion of their nutrients and it would be irresponsible to recommend extensive use of fire in Eastern Canada without investigating this matter.

I am pleased, therefore, to report that both the Research Division of the Ontario Department of Lands and Forests and the Federal Forestry Branch have initiated experiments in prescribed burning in widely different forest cover types. The Ontario Research Division is taking the rather intriguing approach of trying to use fire to control one broadleaved species and favor another. The forest floor under mature stands of sugar maple and yellow birch is being burned in the hope of killing the tangle of sugar maple seedlings and preparing the seed bed for regeneration of yellow birch. This work is being done at Swan Lake in the Algonquin Park. The soil is a deep moulded till with a large nutrient capital and should not suffer greatly from the effects of a fire. The Forestry Branch is making its experiments at the Petawawa Forest Experiment Station and the few trials made to date have been performed in stands of red and white pine growing on a rather dry outwash soil. So far the emphasis has been upon devising appropriate techniques of starting and controlling the burn rather than evaluating its silvicultural effects.

LITERATURE CITED

NEWFOUNDLAND RESEARCH COMMITTEE. 1958. Symposium on prescribed burning in forestry, agriculture and wild life management. Newfoundland Res. Comm. Pub. No. 1. 78 pp.